

Strategik Mineralised Combo Drench for Cattle

Warnings: Harmful if swallowed. May cause an allergic skin reaction. Suspected of causing genetic defects and may damage fertility or the unborn child. May cause damage to organs. Harmful to terrestrial vertebrates.

Handling precautions: Wear protective gloves and use personal protective equipment as required. Avoid breathing vapours. Do not eat, drink or smoke when using this product. Wash hands thoroughly after handling. Wash contaminated clothing and/or protective equipment before reuse.

First Aid: IF SWALLOWED: Rinse mouth. IF ON SKIN: Remove contaminated clothing and wash with plenty of soap and water. IF exposed or concerned, contact the National Poisons centre 0800 POISON (0800 764 766) or a doctor immediately.

Disposal: Avoid release to the environment. Triple rinse empty container, crush or puncture and bury in a suitable landfill, or if appropriate, recycle.

Safety Data Sheet: If additional hazard information is required refer to the Safety Data Sheet. For a copy phone: 0800 587 696.

EFFICACY SPECTRUM

STRATEGIK Mineralised Combo Drench for Cattle is a combination anthelmintic containing a member of both benzimidazoles (white) and imidazothiazoles (clear) drench families. It is effective against parasites sensitive to one or both of these families.

STRATEGIK Mineralised Combo Drench for Cattle is highly effective against the following:

Tapeworms

Mature and immature gastrointestinal roundworms (including those that may be resistant to benzimidazole or levamisole).

Haemonchus, Ostertagia, Trichostrongylus, Nematodirus, Cooperia, Strongyloides, Bunostomum, Oesophagostomum, Chabertia, Trichuris.

Lungworms

Adult Liver Fluke

Resistance can develop to any anthelmintic. Ask your local veterinary

practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

DIRECTIONS FOR USE

SHAKE WELL BEFORE USE

Draft animals into lines of similar bodyweight. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on the heaviest animal in each group (bulls, cows, steers, calves etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Correct drenching technique should be used. Check the accuracy of drench guns regularly.

Dose Rates: Cattle

1 mL/10 kg liveweight (10 mg/kg albendazole and 7.5 mg/kg levamisole HCl, 0.09 mg/kg selenium, 0.2 mg/kg copper, 0.06 mg/kg zinc, 0.02 mg/kg cobalt and 0.09 mg/kg iodine).

DOSE VOLUME TABLE

Bodyweight (kg)	Dose Volume (mL)	10 L treats (head)
60	6	1666
120	12	833
180	18	555
240	24	416
300	30	333
360	36	278
420	42	238
480	48	208
540	54	185
600	60	166
660	66	151

Cattle in excess of 660 kg add 1 mL per 10 kg bodyweight.

Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian.

Do not exceed the stated dose, or dose more frequently than 3-weekly.

Doses of 3 or more times recommended dose can cause symptoms of levamisole toxicity, so estimate liveweights carefully.

Dehydrated animals may be more susceptible to toxicity.

Fatal interactions may occur between levamisole and organophosphate dips.

WITHHOLDING PERIODS:

Meat: Animals producing meat or offal for human consumption must not be sold for slaughter during or within 10 days of cessation of the last treatment.

Milk: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

ERMA Approval No. HSR05082.

Registered pursuant to the ACVM Act 1997, No. A9730 see www.nzfsa.govt.nz/acvm for registration conditions.


DIRECTIONS FOR STORAGE: Store in the original container, tightly closed in a safe place. Store below 30°C (room temperature).



EXP.

Jurox

Registered to Jurox New Zealand Limited
8 Kordel Place
East Tamaki, Auckland
Phone 0800 587 696



409286