

Cobalex Range



1000 x 500ml

Vitamin B12 deficiencies

Before treatment, establish vitamin B12 status of stock and/or pasture cobalt levels. Cobalt deficiency in soil and/or pasture leads to a lack of vitamin B12 in body tissues of stock which subsequently may have poor growth rates and ill thrift, particularly in young animals. Treatment of pregnant animals should benefit newborn animals particularly when treated in the latter stages of pregnancy.

Supplementation of young ruminants should begin at or before weaning. Seek veterinary advise if poor stock condition persists after treatment.

DIRECTIONS FOR USE: Administer by subcutaneous or intramuscular injection, into the anterior half of the neck.

Sheep 1 - 2mL Cattle 4 - 12mL

WITHHOLDING PERIOD: NIL

1000 x 500ml + Se

Selenium and vitamin B12 deficiencies

Before treatment, establish selenium and vitamin B12 status of stock and/or pasture cobalt levels. Do not administer COBALEX PLUS SELENIUM INJECTION unless deficiencies of both cobalt and selenium exist. Cobalt deficiency in soil and/or pasture leads to a lack of vitamin B12 in body tissues of stock. Vitamin B12 deficiency is mostly a problem in very young stock. Poor growth rates and ill thrift may be seen in affected animals. Treatment of pregnant animals should benefit newborn animals particularly when treated in the latter stages of pregnancy.

Supplementation of young ruminants should begin at or before weaning. The need for, and frequency of dosing, dosing is related to the severity of deficiency and the age of the animal. Seek veterinary advise if poor stock condition persists after treatment.

Contraindications: Do not use unless a selenium deficiency has been diagnosed. Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian.

DIRECTIONS FOR USE: Administer by subcutaneous (under the skin) injection into the anterior half of the neck. Avoid intramuscular injection. Use clean sharp needles and replace them frequently. Short needles are recommended.

Lambs: 1mL Adult Sheep 2 - 3mL

Calves 4 - 6 mL Adult Cattle 8 - 12mL

As selenium is toxic above the recommended rates, do not exceed stated dose, or dose more frequently than once every 3 months.

WITHHOLDING PERIOD: NIL