



Presentation

NOROMECTIN Injection is a ready to use solution of ivermectin 10mg/mL, a broad-spectrum antiparasitic for cattle. It is supplied in HDPE vials of 50mL, and 500mL

Indications

NOROMECTIN Injection is effective in the treatment and control of the following

Gastrointestinal roundworms, including 4th stage and – “L4” and adults (including arrested or hypobiotic stages*): *Ostertagia circumcincta**, *O. lyrata*, *Haemonchus spp**, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia oncophora**, *C. punctata**, *Oesophagostomum radiatum*, *Nematodirus spathiger*, *N. helvetianus*, *Strongyloides papillosus*, *Chabertia ovina* and *Tri churls spp.*

Lungworms (immature and adult): *Dictyocaulus viviparus*

Lice: *Linognathus vituli*, *Haematopinus erysternus*, *Solenopotes capillatus*. Aids in control of biting lice, *Damalinia bovis*

Mites *Sarcoptes scabiei* var. *bovis*. Aids in control of *Chorioptes bovis*

Persistent Activity: NOROMECTIN Injection given at the recommended dose rate effectively controls infections with *Ostertagia spp.* acquired at least 14 days after treatment; *Cooperia spp.* acquired up to at least 7 days after treatment; and *Dictyocaulus viviparus* and *Oesophagostomum radiatum* acquired up to at least 21 days after treatment

Administration

NOROMECTIN Injection is to be given to cattle by subcutaneous injection only, through standard injection equipment

Dose Rate

The dose rate is 300µg ivermectin per kg bodyweight; which is provided by 1mL Noromectin Injection for Cattle and Pigs per 33 kg bodyweight. The following table indicates the recommended dose volumes for various bodyweights and the number of doses/pack. Especially in pigs under 16 kg for which less than 0.5 mL Noromectin Injection for Cattle and Pigs is indicated, dosing accuracy is important. The use of a syringe that can be accurate in delivering 0.1 mL is recommended. For piglets weighing less than 16kg, give as 0.1 mL per 3kg.

Bodyweight (kg)	Dose Volume (mL)	Doses Per Pack
33	1.0	200
50	1.5	133
66	2.0	100
99	3.0	66
133	4.0	50
166	5.0	40
200	6.0	33

Over 200 kg bodyweight, give 1.0 mL per 33 kg bodyweight

Withholding Periods:

Meat

Pigs producing meat and offal for human consumption must not be slaughtered within 49 days after treatment



Safe, Guaranteed & On Time

Precautions:

Do not smoke or eat while handling the product. Wash hands after handling the product. If accidentally swallowed, contact a doctor or Poisons Information Centre. In the case of spillage, rinse from eyes or skin with copious amounts of water.

Pharmaceutical Precautions

This product does not contain antimicrobial preservatives. Store below 25°C. Protect from direct sunlight.

Additional Information

Transitory discomfort has been noted following treatment in some cattle. A low incidence of soft tissue swelling at the injection site has been observed. These reactions disappear without treatment

Disposal:

It has been established that ivermectin is readily and tightly bound to soil on contact and is then subject to degradation with time. Drug containers and any residual contents should be disposed of crushing and burying, or incinerating. Avoid contamination of waterways, as free ivermectin may adversely affect fish and some water-borne organisms.

KEEP OUT OF REACH CHILDREN