

POURACIDE NF

CLAIM AND EFFICACY

Pouracide NF is a single dose back-line spray for the control of lice on cattle for up to six weeks.

Pouracide NF also provides control over the irritating effects of biting flies for up to three weeks.

EFFICACY SPECTRUM

Sucking Lice *Linognathus vituli*

WARNING

This substance may be harmful if swallowed or if skin contact occurs.

PRECAUTIONS

- This material may cause irritations in persons with sensitive skin.
- Splashes should be washed from the skin immediately with soap and water.
- Do not contaminate drains, rivers or streams with this product or the container.
- Store product in original container away from food and children.

FIRST AID

If accidentally swallowed, drink water, induce vomiting and call a Doctor immediately.

CAUTION

In rare cases some cows are irritated by synthetic pyrethroids for 48-72 hours after application. This tends to be on a herd basis and repeatable, so do not use on cows known to have been irritated by synthetic pyrethroids previously.

DIRECTIONS FOR USE

- Pouracide NF is a single application back-line spray for cattle.
- Wear gloves when treating cattle.
Apply Pouracide NF to the back from the withers to the base of the tail by means of calibrated applicator or Protector Applicator gun.
- Store below 30°C.
- Destroy empty containers.

DOSE RATES

Example : 10mL per 100 kg bodyweight

Bodyweight	Dose Volume	5L Treats
100 kg	10 mL	500
200 kg	20 mL	250
300 kg	30 mL	166
400 kg	40 mL	125

WITHHOLDING PERIOD

MILK: Nil

MEAT: Cattle producing meat and offal for human consumption must not be slaughtered during or within 7 days of the last treatment.

Jurox

Licensed under the Animal Remedies Act 1967, No 6438
Licensed to Jurox New Zealand Limited
1770 Ponga Road, R. D. Papakura,
Auckland
Ph: 0800 587 696