

## Promectin

### INDICATIONS

**PROMECTIN PLUS Allwormer Paste for Horses** is highly effective against tapeworms and gastrointestinal, cutaneous and pulmonary nematodes and bots of horses of all ages. Regular treatment will reduce the chances of verminous arteritis and colic caused by *Strongylus vulgaris*. At the recommended dose level of 1mL per 20kg bodyweight, **PROMECTIN PLUS Allwormer Paste for Horses** controls the following parasites:

**Tapeworms:** *Anoplocephala perfoliata*, *Anoplocephala magna*, *Paranoplocephala mammilana* (adult and immature, heads and segments).

**Large Strongyles:** *Strongylus vulgaris* (adults and arterial stages), *Strongylus edentatus* (adults and tissue stages), *Strongylus equinus* (adults) and *Triodontophorus* spp (adults).

**Small Strongyles:** including benzimidazole resistant strains (adult and immature); *Cyathostomum* spp, *Cylicocyclus* spp, *Cylicostephanus* spp, *Cylicodontophorus* spp, *Gyalocephalus* spp.

**Pinworms:** *Oxyuris equi* (adult and immature).

**Ascarids:** *Parascaris equorum* (adult and immature).

**Hairworms:** *Trichostrongylus axei* (adult).

**Large Mouth Stomach Worms:** *Habronema muscae* (adult).

**Neck Threadworms:** *Onchocerca* spp (microfilariae).

**Bots:** *Gasterophilus* spp (oral and gastric stages).

**Lungworms:** *Dictyoacaulus arnfieldi* (adult and immature).

**Intestinal Thread Worms:** *Strongloides westeri* (adult).

**PROMECTIN PLUS Allwormer Paste for Horses** also effectively controls skin lesions caused by *Habronema* and *Draschia* spp, cutaneous larvae (summer sores) and microfilariae of *Onchocerca* spp (cutaneous onchocerciasis). All horses should be included in a regular parasite control programme, with particular attention paid to mares, foals and yearlings. Because re-infection is common, treatment must be repeated routinely for effective control. Contact your veterinarian for a control programme to meet your needs.

### DIRECTIONS FOR USE

**Dose:** Dose orally at the recommended rate of 0.2mg/kg abamectin and 2.5mg/kg praziquantel. Each weight marking on the syringe plunger will deliver sufficient paste to treat 50kg bodyweight. The contents of the syringe will treat a total 600kg bodyweight.

**Administration:** Turn the ring on the plunger to the appropriate bodyweight setting. Insert the syringe into the side of the horse's mouth, in the gap between the front and the back teeth. Advance the plunger as far as it will go, depositing the paste on the rear of the tongue. Immediately raise the horse's head for a few seconds after dosing, to ensure the full dose has been swallowed.

### MEAT WITHHOLDING PERIOD

**DO NOT USE less than 28 days before slaughter for human consumption.**