

### Strategik Combo

Dual Action Mineralised Sheep and Lamb Drench



Strategik Combo Dual Action Mineralised Sheep and Lamb Drench is intended for use against benzimidazole and levamisole susceptible worms of the following species:

**ROUNDWORMS** (mature and immature)

Barber's pole worm	Haemonchus contortus
Small brown stomach worm	Ostertagia spp.
Stomach hair worm	Trichostrongylus axei
Black sour worm	Trichostrongylus spp.
Small intestinal worm	Cooperia spp.
Hookworm	Bunostomum spp.
Nodule worm	Oesophagostomum columbianum
Thin-necked intestinal worm	Nematodirus spp.
Large bowel worm	Oesophagostomum venulosum
Large-mouthed bowel worm	Chabertia ovina

**LUNGWORM**

Large lungworm	Dictyocaulus filaria
----------------	----------------------

**TAPEWORM**

Heads and Segments	Moniezia spp.
--------------------	---------------

**LIVER FLUKE** (Adults)

Fasciola hepatica
-------------------

**OVICIDAL**: Reduces the output of viable worm and fluke eggs.

Strategik Combo also assists in correction of copper, cobalt and selenium deficiency.

Strategik Combo contains both albendazole, a member of the benzimidazole family and levamisole. It is effective against internal parasites sensitive to one or both of these families. Resistance can develop to any anthelmintic. Check the effectiveness of anthelmintics at least annually. Follow recommended parasite management practices to reduce the potential for resistance to develop. Alternate between anthelmintic families, ideally on a yearly basis. Do not alternate between drench families within the same season.

**Dose Rates**

Sheep and Lambs	1mL per 5kg body weight (4.75 mg/kg albendazole 7.5 mg/kg levamisole)
-----------------	---

**Dose Volume Table**

Body Weight (kg)	Dose Volume (mL)	20 Litres Treats (head)
10	2	10,000
15	3	6,666
20	4	5,000
25	5	4,000
30	6	3,333
35	7	2,875
40	8	2,500
45	9	2,222
50	10	2,000
55	11	1,818
60	12	1,666

Sheep in excess of weights above to be dosed at the appropriate rate shown before the Dose Volume Table.

**Withholding periods:**

**Animals producing meat and offal for human consumption must not be slaughtered during or within 10 days of the last treatment. Milk intended for human consumption must be discarded during treatment and for 10 days following the last treatment.**

Licensed under the Animal Remedies Act 1967, No 7700  
 Licensed to Jurox New Zealand Limited  
 1770 Ponga Road, Papakura  
 Phone 0800 587 696.