

Combination Pour-on for Cattle BZ/Lev

ACTIVE INGREDIENTS: Oxfendazole 150mg/mL, and Levamisole 100mg/mL

INDICATIONS

A broad spectrum combination (containing oxfendazole and levamisole) pour on anthelmintic for the control of susceptible gastrointestinal roundworms and lungworms in cattle including benzimidazole and levamisole resistant strains of the following internal parasites:

Gastrointestinal roundworms: *Ostertagia ostertagi*, *Ostertagia leptospicularis*, *Ostertagia lyrata*, *Teladorsagia (Ostertagia) circumcincta*, *Haemonchus contortus*, *Haemonchus placei*, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Trichostrongylus vitrinus*, *Trichostrongylus longispicularis*, *Cooperia oncophora*, *Cooperia punctata*, *Cooperia curticei*, *Strongyloides papillosus*, *Nematodirus helvetianus*, *Nematodirus spathiger*, *Bunostomum phlebotomum*, *Oesophagostomum radiatum*, *Oesophagostomum venulosum*, *Trichuris ovis*

Lungworm: *Dictyocaulus viviparus*

Tapeworm: *Moniezia expansa*

Also for the treatment of Type 11 *Ostertagia* and will reduce the number of viable eggs produced by all the important gastrointestinal roundworms

DIRECTIONS FOR USE

Shake well before use

Apply topically along the backline beginning at the withers and ending at the base of the tail
Combination Pour On for Cattle is a ready to use anthelmintic that can be dosed through standard pour on application equipment.

It is important to give the correct dose.

Before treatment, weigh a representative sample of cattle to determine the weight of the heaviest animal in each group

Use correct dosing techniques and check accuracy of applicator regularly

It is advised that a resistance test be carried out regularly when using any parasite treatment

DOSE RATE: 1ml per 10kg bodyweight

Bodyweight	Dose Volume
60-70kg	7mL
71-80kg	8mL
91-100kg	10mL
101-120kg	12mL
141-160kg	16mL
181-200kg	20mL
241-260kg	26mL
281-300kg	30mL

WITHHOLDING PERIODS:

MEAT: Cattle producing meat or offal for human consumption must not be sold for slaughter during or within 35 days of cessation of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment

Registered pursuant to the ACVM Act 1997, No. A10326