

Combination Sheep Drench

Combination Sheep Drench 20Litre

A Broad Spectrum Combination sheep drench for the control of susceptible gastrointestinal roundworms and lungworms in sheep and lambs including benzimidazole and levamisole resistant strains

INDICATIONS

Contains albendazole (23.8 mg/mL), levamisole HCl 37.5 mg/mL, selenium 0.5 mg/mL, cobalt 0.3 mg/mL, copper 2.1 mg/mL, zinc 0.6 mg/mL.

FOR THE TREATMENT OF: tapeworms and liver fluke and the following susceptible adult and immature stages of gastrointestinal roundworms and lungworms including benzimidazole and levamisole resistant strains: *Haemonchus contortus*, *Ostertagia* spp., *Trichostrongylus* spp., *Nematodirus* spp., *Cooperia* spp., *Strongyloides papillosus*, *Bunostomum trigonocephalum*, *Chabertia ovina*, *Oesophagostomum venulosum*, *Trichuris ovis* (adults only), *Moniezia expansa* (tapeworm segments only), *Dictyocaulus* spp., *Muellerius capillaris*, Liver fluke (adults only). Also kills roundworm and fluke eggs.

DIRECTIONS FOR USE

Shake well before use. Combination Sheep Drench is a ready to use anthelmintic that can be dosed through standard drenching equipment. It is important to give the correct dose.

DOSE RATE: 1mL/5kg

16-20kg - 4mL -	5000 doses per 20Litre
26-30kg - 6mL -	3332 doses per 20Litre
36-40kg - 8mL -	2500 doses per 20Litre
46-50kg - 10mL-	2000 doses per 20Litre
66-70kg - 14mL -	1428 doses per 20Litre

WITHHOLDING PERIODS:

MEAT - Sheep producing meat or offal for sale for human consumption must not be sold for slaughter either

During treatment or within 10 days of the last treatment

MILK – Milk intended for sale for human consumption must be discarded during treatment and for nor less than 35 days following last treatment

Registered pursuant to the ACVM Act 1997, No. A9764.