

Noromectin Injection for Sheep

Noromectin Injection for Cattle, Sheep & Pigs is a ready to use solution of ivermectin 10mg/mL, a broad-spectrum antiparasitic treatment for cattle, sheep & pigs
It is supplied in HDPE flexible packs 500mL

SHEEP: Noromectin Injection for Cattle, Sheep & Pigs is effective in the treatment and control of the following –

Gastrointestinal roundworms: *Haemonchus contortus* (adults, L4, L3). *Ostertagia circumcincta* (adults, L4, L3, including inhibited larvae). *O. trifurcata* (adults, L4) *Trichostrongylus axei*, (adults). *T. colubriformis* (adults, L4 L3) *T. vitrinus* (adults). *N. spathiger* (L3). *Cooperia curticei* (adults L3) *O. venulosum* (adults). *Chabertia ovina* (adults, L4, L3). *Strongyloides papillosus* (L4). **Lungworms:** *Dictyocaulus filaria* (adults, L4, L3)

Nasal Bot: (all larval stages) *Oestrus ovis*.

Itch Mite: *Psorergates ovis*

ADMINISTRATION:

Noromectin Injection for Cattle, Sheep & Pigs is to be given to cattle, sheep and pigs by subcutaneous injection only. Standard injection equipment is suitable for administration.

DOSE RATE: 0.1mL per 5kg bodyweight

| | | |
|------------|-------|----------------------|
| 11 - 15 kg | 0.3mL | 1666 doses per 500ml |
| 16 - 20 kg | 0.4mL | 1250 doses per 500ml |
| 21 - 25 kg | 0.5mL | 1000 doses per 500ml |
| 26 - 30 kg | 0.6mL | 833 doses per 500ml |
| 46 - 50 kg | 1.0mL | 500 doses per 500ml |
| 61 - 70 kg | 1.4mL | 357 doses per 500ml |
| 71 - 80 kg | 1.6mL | 312 doses per 500ml |

WITHHOLDING PERIODS:

MILK: Cows & Sheep: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

MEAT: Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 35 days of the last treatment.

PRECAUTIONS:

This product is not for intravenous or intramuscular use. Noromectin Injection for Cattle, Sheep & Pigs is a low-volume product registered for use in the stated species. It should not be used in other species as severe adverse reactions, including fatalities in dogs, may occur.

Discard contents unused after 6 months from broaching the pack. Store below 25°C. Protect from direct sunlight.

Registered pursuant to the ACVM Act 1997 No. A8208