

Noromectin Oral + Se Liquid for Sheep

Noromectin Oral Plus Selenium is a ready to use solution of Ivermectin 0.8mg/mL with 0.4mg/mL Selenium, as sodium selenate for the use as a broad-spectrum antiparasitic treatment for sheep

INDICATIONS:

Noromectin Oral Plus Selenium is effective in the treatment and control of the following

Gastrointestinal roundworms: including infective 3rd stage –‘L3’, 4th stage –‘L4’ and adults (including arrested or hypoiotic stages*):

Haemonchus contortus* Ostertagia circumcincta* Trichostrongylus axei, T colubriformis T vitrinus Cooperia oncophora, C curticei, Nematodirus. Spathiger, N filicollis,) Oesophagostomum venulosum. Strongyloides papillosus, Chabertia ovina and Trichuris ovis

Lungworms: (immature and mature) Dictyocaulus filaria

Nasal Bot: (all larval stages) Oestrus ovis.

Itch Mite: Psorergates ovis

ADMINISTRATION: Noromectin Oral Plus Selenium is given orally through a standard drench gun, it has a wide safety margin and may be used at the recommended dose in sheep of all ages, and in breeding animals

DOSE RATE: 2.5mL per 10kg of bodyweight

Weight Range	Dose Volume	Doses per 20 Litre
11-20kg	5.0ml	4000
21-30kg	7.5ml	2666
31-40kg	10.0ml	2000
41-50kg	12.5ml	1600
51-60kg	17.5ml	1142
71-80kg	20.0ml	1000

Resistance can develop to any anthelmintic.. To delay the onset of resistance:

- * It is advisable that a resistance test be conducted regularly when using any parasite treatment
- * Before dosing, weigh a group of animals to accurately calculate the dose
- * Choose the dose according to the heaviest animal
- * Do not under-dose
- * Check the accuracy of drench guns regularly
- * Do not change between drench families within the same season
- * Ask your local veterinarian or animal health adviser about recommended parasite management practices for your area. Follow these recommendations to reduce the potential for resistance to develop.

WITHHOLDING PERIODS:

Meat: Sheep producing meat and offal for human consumption must not be slaughtered within 10 days after treatment.

Milk: Do not treat sheep producing milk for human consumption, or within 28 days before beginning lactation for human consumption.

Registered pursuant to the ACVM Act 1997 No. A9449