

## Noromectin Pour-on for Deer

Noromectin Pour-on for Deer & Cattle is a ready to use solution containing Ivermectin 5mg/mL, a broad-spectrum antiparasitic treatment for cattle and deer

### INDICATIONS:

Deer: Indicated for the treatment and control of parasites sensitive to Ivermectin; including the following- **Gastrointestinal roundworms**, adults Ostertagia types – Ostertagia leptospicularis, Spiculopteragia asymmetrica, Skrjabinagia kolchida and adult stages of Oesophagostomum sp

**Lungworms** (adult and fourth stage larvae): Dictyocaulus viviparus

**Persistent Activity:** Given at the recommended dose rate, Noromectin Pour-On effectively controls infections with Dictyocaulus viviparus acquired up to 28 days after treatment.

### ADMINISTRATION:

NOROMECTIN Pour-On is a ready to use solution. Apply topically along the midline of the back from the withers to the base of the tail.

**DOSE RATE:** 1ml per 10kg bodyweight

<b>Weight Range</b>	<b>Dose Volume</b>
Up to 100kg	Up to 10ml
101-150kg	10-15ml
151-200kg	15-20ml
201-250kg	20-25ml
251-300kg	25-30ml
301-350kg	30-35ml
351-400kg	35-40ml
401-450kg	40-45ml
451-500kg	45-50ml
501-550kg	50-55ml
551-600kg	55-60ml
601-650kg	60-65ml

Noromectin Pour-on for Cattle and Deer has a wide safety margin and may be used at the recommended dose in cattle and deer of all ages and in breeding animals.

### WITHHOLDING PERIODS

**Meat** -Deer producing meat and offal for human consumption must not be slaughtered within 21days after treatment

Registered pursuant to the ACVM Act 1997 No 8032