

Q-drench multicomination drench for Sheep

Q-drench Multi-Combination Drench for Sheep

Active Constituents: Abamectin 1.0 g/L, Albendazole 25.0 g/L, Closantel 37.5 g/L, Levamisole hydrochloride 40.0 g/L

~Quarantine drench~

There is no known resistance to Q-Drench in NZ

For the treatment and control in sheep of susceptible gastro-intestinal roundworms It is also effective against lungworm, tapeworms, mature and late immature liver fluke, nasal bot and itch mite.

Q-drench is a broad-spectrum oral anthelmintic which can be used against susceptible gastrointestinal parasites and to prevent the introduction of resistant strains onto a property.

INDICATIONS

Adult and immature gastrointestinal roundworms:

Barber's Pole Worm *Haemonchus contortus* - including inhibited L4 stage and strains with single, dual or triple resistance to macrocyclic lactones, benzimidazoles, levamisole or closantel, and strains with emerging resistance to closantel.

Small Brown Stomach Worm *Teladorsagia (Ostertagia) circumcincta*

- including inhibited L4 stage and strains with single or dual resistance to macrocyclic lactones, benzimidazoles or levamisole:

Black Scour Worm

Trichostrongylus colubriformis - including strains with single or dual resistance to macrocyclic lactones, benzimidazoles or levamisole.

- * Large Stomach Worm *Haemonchus placei*
- * Stomach Hair Worm *Trichostrongylus axei*
- * Black Scour Worm *Trichostrongylus vitrinus*
- * Small Intestinal Worm *Cooperia* spp
- * Thin-Necked Intestinal Worm *Nematodirus* spp
- * Large Mouthed Bowel Worm *Chabertia ovina*
- * Nodule Worm *Oesophagostomum columbianum*
- * Large Bowel Worm *Oesophagostomum venulosum*
- * Whip worm *Trichuris ovis*
- * Intestinal threadworm *Strongyloides papillosus*
- * Hookworm *Bunostomum* spp

Nasal Bot *Oestrus ovis*

Lungworm *Dictyocaulus viviparus* (Large lungworm)

Liver Fluke *Fasciola hepatica* - mature fluke and immature forms including 6 week old stages. Reduces the output of viable fluke eggs.

Tapeworms *Monezia* spp - heads and segments

Itch mite *Psorergates ovis*

DIRECTIONS FOR USE

For use as a quarantine drench: Treat all animals on arrival, or ideally at farm of origin, and hold in yards for 24 hours before release. For use in a strategic worm control program: Speak to your animal health adviser or veterinarian. Administer orally

DOSE RATE 1ml per 5kg bodyweight

| Live weight | Dose Volume | 10L treats |
|-------------|-------------|------------|
| 21 - 25 kg | 5ml | 2000 |
| 26 - 30 kg | 6ml | 1666 |
| 31 - 35 kg | 7ml | 1428 |
| 36 - 40 kg | 8ml | 1250 |
| 41 - 45 kg | 9ml | 1111 |



14 Athenaeum Rd,
RD1, Craigellachie
Lawrence
Free Phone 0800 4 DRENCH (437 3624)
Phone/Fax ++64 3 485 9427
Email call-us@drench.co.nz

| | | |
|------------|------|------|
| 46 - 50 kg | 10ml | 1000 |
| 51 - 55 kg | 11ml | 909 |
| 56 - 60 kg | 12ml | 833 |
| 61 - 65 kg | 13ml | 769 |
| 66 - 70 kg | 14ml | 714 |
| 71 - 75 kg | 15ml | 666 |

Shake well before use

WITHHOLDING PERIODS

MEAT - Do NOT use less than 28 days before slaughter for human consumption.

MILK - Do NOT use in female sheep which are producing or may in the future produce milk or milk products for human consumption.

Registered pursuant to the ACVM Act 1997, No. A9291