

Strategik Cattle Combination

Albendazole Levamisole

ACTIVE CONSTITUENTS: 100g/L Albendazole, 75g/L Levamisole, 0.2g/L Cobalt 2.0g/L Copper, 0.6g/L zinc, 0.9g/L iodine, and 0.9g/L Selenium

EFFICACY SPECTRUM

STRATEGIK Mineralised Combo Drench for Cattle is a combination anthelmintic containing a member of both benzimidazoles (white) and imidazothiazoles (clear) drench families. It is effective against parasites sensitive to one or both of these families.

INDICATIONS:

STRATEGIK Mineralised Combo Drench for Cattle is effective against the following:

Tapeworms

Mature and immature gastrointestinal roundworms (including those that may be resistant to benzimidazole or levamisole). *Haemonchus*, *Ostertagia*, *Trichostrongylus*, *Nematodirus*, *Cooperia*, *Strongylus*, *Bunostomum*, *Oesophagostomum*, *Chobertia*, *Trichuris*. Lungworm

Lungworm

Adult Liver Fluke

DIRECTIONS FOR USE

Shake well before use

Draft animals into lines of similar bodyweight. A representative sample of animals should be weighed before treatment either with scales or a weighband.

Dose rate to be based on the heaviest animal in each group (bulls, cows, steers, calves etc.).

Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Correct drenching technique should be used. Check the accuracy of drench guns regularly.

DOSE RATE: 1ml per 10kg bodyweight

Bodyweight	Dose Volume	Doses per 10Lt
60kg	6mL	1666
120kg	12mL	833
180kg	18mL	555
240kg	24mL	416
300kg	30mL	333
360kg	36mL	278
420kg	42mL	238
480kg	48mL	208

PRECAUTIONS Do not exceed the stated dose, or dose more frequently than 3-weekly.

Doses of 3 or more times recommended dose can cause symptoms of levamisole toxicity, so estimate live weights carefully.

Do not use at the same time as any other selenised fertilizer, prill or product without consulting a veterinarian. Dehydrated animals may be more susceptible to toxicity.

Fatal interactions may occur between levamisole and organophosphate dips.

WITHHOLDING PERIODS:

MEAT: Animals producing meat or offal for human consumption must not be sold for slaughter during or within 10 days of cessation of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment,



14 Athenaeum Rd,
RD1, Craigellachie
Lawrence
Free Phone 0800 4 DRENCH (437 3624)
Phone/Fax ++64 3 485 9427
Email call-us@drench.co.nz