

Troika - Triple-active sheep drench

ACTIVE CONSTITUENTS: 1.0g/L Abamectin, 25g/L Albendazole and 40.0g/L Levamisole hydrochloride

INDICATIONS: TROIKA Combination Drench for Sheep contains abamectin, albendazole and levamisole, which are members of the macrocyclic lactone, benzimidazole and imidazothiazole groups respectively. It is a broad-spectrum oral anthelmintic which can be used against susceptible gastrointestinal parasites and to prevent the introduction of resistant strains on to a property.

It is effective against susceptible strains of the following parasites of sheep:

Adult and immature gastrointestinal roundworms:

Barber's Pole Worm (*Haemonchus contortus*) - including inhibited L4 stage. Small Brown Stomach Worm (*Teladorsagia (Ostertagia) circumcincta*) - including inhibited L4 stage. Black Scour Worm (*Trichostrongylus colubriformis*), Large Stomach Worm (*Naemonchus placei*)

Stomach Hair Worm (*Trichostrongylus axei*), Black Scour Worm (*Trichostrongylus vitrinus*)

Small Intestinal Worm (*Cooperia* spp), Thin-Necked Intestinal Worm (*Nematodirus* spp)

Large Mouthed Bowel Worm (*Chabertia ovina*), Nodule Worm (*Oesophagostomum columbianum*)

Large Bowel Worm (*Oesophagostomum venulosum*), Whip worm (*Trichuris ovis*)

Intestinal threadworm (*Strongyloides papillosus*), Hookworm (*Bunostomum* spp)

Lungworm - Large lungworm (*Dictyoxia viviparus*)

Tapeworms (*Monieria* spp.)

Liver Fluke –adults (*Fasciola hepatica*) reduces the output of viable fluke eggs

Itch Mite (*Psorergates ovis*)

Nasal Bot (*Oestrus Ovis*)

DIRECTIONS FOR USE:

Shake well before use

Administer Orally and check accuracy of the drench equipment before and during use

Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose

Doses of 3 or more times recommended dose can cause symptoms of levamisole toxicity, so estimate liveweights carefully.

Dehydrated animals may be more susceptible to toxicity.

Fatal interactions may occur between levamisole and organophosphate dips.

DOSE RATE: 1ml per 5kg bodyweight

Bodyweight	Dose Volume	Doses per 10Lt
21-25kg 5mL	3000	
26-30kg 6mL	2500	
31-35kg 7mL	2142	
36-40kg 8mL	1875	
46-50kg 10mL	1500	
56-60kg 12mL	1250	
66-70kg 14mL	1071	
71-75kg 15mL	1000	

WITHHOLDING PERIODS:

MEAT: Animals producing meat or offal for human consumption must not be sold for slaughter during or within 14 days of cessation of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment,